16:28

Application Serial No. 10/715,773
Preliminary Amendment dated September 16, 2005

Amendments to the Claims:

This listing of claims will replace all prior versions and listing of claims in the application.

Listing of Claims:

1. (Original) A compound of formula I:

$$R_1$$
 R_2
 R_3
 R_4
 R_{10}
 R_{10}
 R_{10}
 R_{10}
 R_{10}

wherein:

X is HN, R₁₁N, S, O, CH₂, or R₁₁CH;

R₁₁ is hydrogen, (C₁-C₄)alkyl, or (C₁-C₄)alkanoyl;

 R_1 - R_8 are each independently hydrogen, hydroxy, mercapto, amino, nitro, $(C_1$ - $C_4)$ alkyl, $(C_1$ - $C_4)$ alkoxy, $(C_1$ - $C_4)$ alkylthio, or halo; wherein two adjacent groups of R_1 - R_5 together with the phenyl ring to which they are attached may optionally form a fused ring, for example forming a naphthyl or a tetrahydronaphthyl ring; and further wherein the ring formed by the two adjacent groups of R_1 - R_5 may optionally be substituted by 1, 2, 3, or 4 hydroxy, mercapto, amino, nitro, $(C_1$ - $C_4)$ alkyl, $(C_1$ - $C_4)$ alkoxy, $(C_1$ - $C_4)$ alkylthio, or halo; and

 R_9 and R_{10} are each independently hydrogen, (C_1 - C_4)alkyl, (C_1 - C_4)alkoxy, halo, or (C_1 - C_4)alkanoyl; or R_9 and R_{10} together are methylenedioxy; or a pharmaceutically acceptable salt thereof;

Page 3 of 9

Application Scrial No. 10/715,773 Preliminary Amendment dated September 16, 2005

FROM-Merchant & Gould

provided the compound is not 4-(4'-hydroxyl-phenyl)-amino-6,7-dimethoxyquinazoline.

(Canceled) 2-12.

13. (Original) A pharmaceutical composition comprising a compound of formula I:

$$R_1$$
 R_2
 R_3
 R_4
 R_{10}
 R_{10}
 R_{10}
 R_{10}
 R_{10}
 R_{10}

wherein:

X is HN, $R_{11}N$, S, O, CH_2 , or $R_{11}CH$;

 R_{11} is hydrogen, (C_1-C_4) alkyl, or (C_1-C_4) alkanoyl;

R₁-R₈ are each independently hydrogen, hydroxy, mercapto, amino, nitro, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, or halo; wherein two adjacent groups of R₁-R₅ together with the phenyl ring to which they are attached may optionally form a fused ring, for example forming a naphthyl or a tetrahydronaphthyl ring; and further wherein the ring formed by the two adjacent groups of R₁-R₅ may optionally be substituted by 1, 2, 3, or 4 hydroxy, mercapto, amino, nitro, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkylthio, or halo; and R_9 and R_{10} are each independently hydrogen, (C_1-C_4) alkyl, (C_1-C_4) alkoxy, halo, or (C₁-C₄)alkanoyl; or R₉ and R₁₀ together are methylenedioxy; or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier.

Page 4 of 9

16:28

Application Serial No. 10/715,773
Preliminary Amendment dated September 16, 2005

14-16. (Canceled)

- 17. (Original) A pharmaceutical composition comprising 4-(4'-hydroxyl-phenyl)-amino-6,7-dimethoxyquinazoline; or a pharmaceutically acceptable salt thereof; and a pharmaceutically acceptable carrier.
- 18. (Original) A therapeutic method for treating leukemia or lymphoma in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 19. (Canceled)
- 20. (Original) A therapeutic method for preventing or reducing ultraviolet B radiation-induced inflammatory response in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 21. (Original) A therapeutic method for inhibiting the release of prostaglandin E₂ in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 22. (Original) A therapeutic method for preventing or reducing UVB-induced skin edema or vascular permeability changes in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 23. (Original) A therapeutic method for preventing or reducing ultraviolet B radiation-induced damage to epithelial cells or mutation frequency in skin in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.

16:28

Application Serial No. 10/715,773
Preliminary Amendment dated September 16, 2005

- 24. (Original) A therapeutic method for protecting a mammal from tumorigenic effects of UVB light comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 25. (Original) A therapeutic method for inhibiting T-cell activity in a mammal comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.
- 26. (Original) A therapeutic method for preventing or treating an autoimmune disease comprising administering to the mammal in need thereof an effective amount of a JAK-3 inhibitor.

27-29. (Canceled)

- 30. (New) A method of treating graft versus host disease comprising administering a JAK-3 inhibitor.
- 31. (New) The method of claim 30, wherein the JAK-3 inhibitor has molecular dimensions compatible with the shape of a JAK-3 kinase binding pocket model and occupies a molecular volume of less than about 530 Å³.
- 32. (New) The method of claim 31, wherein the molecular volume is about 252 $Å^3$ to about 307 $Å^3$.
- 33. (New) The method of claim 32, wherein the JAK-3 inhibitor has an estimated K_{ℓ} of less than about 2.3 μ M as measured by inhibition of JAK-3-induced cell growth.

Application Serial No. 10/715,773
Preliminary Amendment dated September 16, 2005

- 34. (New) The method of claim 33, wherein the estimated K_i is of about 0.6 μ M to about 2.3 μ M.
- 35. (New) The method of claim 31, wherein said JAK-3 inhibitor has the following molecular features:
 - a) a relatively planar molecular shape;
 - b) an ability to fit the 530 Å³ binding pocket volume; and
 - c) a hydroxyl group capable of interacting with Asp 967 of the binding pocket.
- 36. (New) The method of claim 35, wherein said JAK-3 inhibitor further comprises:
 - d) a hydrogen bond acceptor or a hydrogen bond donor capable of interacting with Leu 905 of the binding pocket.
- 37. (New) A method of treating host rejection of a donor organ transplant comprising administering to the host a JAK-3 inhibitor.
- 38. (New) The method of claim 37, wherein the JAK-3 inhibitor has molecular dimensions compatible with the shape of a JAK-3 kinase binding pocket model and occupies a molecular volume of less than about 530 Å³.
- 39. (New) The method of claim 38, wherein the molecular volume is about 252 $Å^3$ to about 307 $Å^3$.
- 40. (New) The method of claim 39, wherein the JAK-3 inhibitor has an estimated K_i of less than about 2.3 μ M as measured by inhibition of JAK-3 positive Leukemia cell growth.

Application Serial No. 10/715,773
Preliminary Amendment dated September 16, 2005

- 41. (New) The method of claim 40, wherein the estimated K_i is of about 0.6 μ M to about 2.3 μ M.
- 42. (New) The method of claim 38, wherein said JAK-3 inhibitor has the following molecular features:
 - a) a relatively planar molecular shape,
 - b) an ability to fit into the 530 Å³ binding pocket volume, and
 - c) a hydroxyl group capable of interacting with Asp 967 of the binding pocket.
- 43. (New) The method of claim 42, wherein said JAK-3 inhibitor further comprises:
 - a hydrogen bond acceptor or a hydrogen bond donor capable of interacting with Leu 905.